#### Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers, Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

### Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers,

Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine

Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

## Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers, Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

## Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers,

Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine

Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

# Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers,

Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine

Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

#### Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers, Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

#### Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers,

Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine

Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

### Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers,

Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine

Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

#### Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers, Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.

# Special Dietary Consideration for 24 Hour Urine Collection for 5-HIAA

Patient should avoid food high in indoles: avocado, banana, tomato, plum, walnut, pineapple, and eggplant.

Drugs which can interfere with testing due to physiologic response: Acetaminophen, Alpha and Beta Blockers,

Atenolol, Bromocriptine, Bronchodilators, Clonidine, Digoxin, Isoniazid, L-Dopa, Labetelol, Methyldopa, Monoamine

Oxidase Inhibitors, Nitroglycerin, Sympathomimetic Amines, Phenobarbital, Phenothiazines, Phentolamine, Reserprine, Salicylates, Tricyclic Antidepressants.